

541,315

10/541315
Rec'd PCT
30 JUN 2005

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
5 August 2004 (05.08.2004)

PCT

(10) International Publication Number
WO 2004/065596 A2

(51) International Patent Classification: C12N 15/00

(21) International Application Number:
PCT/EP2004/000645

(22) International Filing Date: 21 January 2004 (21.01.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03075207.5 21 January 2003 (21.01.2003) EP(71) Applicant (for all designated States except US):
CROPDESIGN N.V. [BE/BE]; Technologiepark 3,
Zwijnaarde 9052 (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HATZFELD, Yves
[FR/FR]; 186 rue des Postes, Lille 59000 (FR). INZE, Dirk
[BE/BE]; Driesstraat 18, Moorsel- Aalst 9310 (BE).(74) Common Representative: CROPDESIGN N.V.; Tech-
nologiepark 3, Zwijnaarde 9052 (BE).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: REGULATORY SEQUENCE

(57) Abstract: The present invention relates to the field of plant molecular biology. In particular, it describes the use of a regu-
latory nucleic acid sequence of the rice gene GOS2 for the regulation of gene expression in plant cells derived from plants other
than monocotyledonous plants. The use of the regulatory sequence of the present invention results in constitutive expression with
expression levels similar to that of CaMV 35S. The present invention also relates to vectors and host cells comprising these nucleic
acid sequences. The invention further relates to transgenic cells and plants comprising these sequences and to methods for obtaining
such cells and plants.

WO 2004/065596 A2

Regulatory sequence

5 The present invention relates to the field of plant molecular biology. In particular, the present invention describes the use of a regulatory sequence of the rice GOS2 gene for the regulation of gene expression in plant cells of non-monocotyledonous plants.

10 Plant genetic engineering has the potential to enable the production of plants having all manner of desirable traits, such as increased pathogen resistance, herbicide resistance, increased yield, increased stress tolerance etc. This is typically achieved by inserting suitable pieces of DNA from a heterologous source, or from the same source, into a plant. The expression of heterologous DNA sequences in a plant host is dependent upon the presence of an operably linked promoter that is functional within the plant host.

15 Promoters are DNA elements that regulate transcription in prokaryotic and eukaryotic cells, and are located in the 5'-flanking or upstream region of a transcribed gene. The sequence of a promoter varies in length and base pair composition from gene to gene. Many regulatory sequence elements may be present within the sequence of a promoter. These regulatory sequence elements, which interact with DNA binding proteins (that are part of the transcription initiation complex) are generally embedded in a promoter sequence. Specific elements, called
20 DNA-boxes, contribute defined features or a defined activation pattern to a promoter. Many of these boxes are recognition sites to which regulatory transcription factors can bind and are part of a tight control mechanism of the promoter. Numerous plant promoter DNA boxes have been described, such as tissue-specific boxes (Chaubet *et al.*, Plant J. 10 (3) 425-435, 1996), pyrimidin boxes (Gubler & Jacobsen, Plant Cell 4 (11) 1435-1441, 1992) which influence the
25 level of expression, or G-boxes (Dolferus *et al.*, Plant Physiol. 105 (4) 1075-1087, 1994), which reduce promoter activity upon exposure to cold or dehydration.

30 Recombinant DNA technology has now found a wide application in research and in product development. In molecular biology, specific types of promoters are frequently used for the expression of heterologous nucleic acids or proteins in organisms, in which the expression may be constitutive (i.e. continuous expression throughout the cells of an organism), or limited to defined parts of an organism (i.e. tissue or cell preferred expression), or limited to certain developmental stages or to certain environmental or physiological conditions (i.e. inducible expression).

35 A need exists in various industrial applications for transcriptional control elements capable of driving gene-expression in plants. Therefore, in order to obtain desired gene expression, which

can be fine tuned depending on specific needs, it is desirable to provide a wide array of promoters from which one can be chosen. Persons skilled in the art need to have at their disposal a variety of promoters with different expression patterns or with different time-control features from which to choose. For many industrial and agronomic applications, where the aim is to confer a given characteristic throughout plant development and during any environmental or physical state of the plant, a constitutive promoter would be needed. A constitutive promoter may be suitable for driving expression of selectable marker genes or may be used when the gene product of a particular gene is to be isolated and purified from the plant for commercial purposes.

It is also frequently desirable to use promoters that are regulated in a comparable way to each other and result in a similar expression pattern over a wide range of host species. Very few constitutive plant promoters that are active in a wide range of species have been isolated to date (Callis *et al.*, J. Biol. Chem. 265 12486-12493, 1990; Zhang *et al.*, Plant Cell 3 1155-1165, 1991; de Pater *et al.*, Plant J. 2 837-844, 1992; Mandel *et al.* Plant Mol. Biol. 29 995-1004, 1995; Baszczyński *et al.*, Maydica 42 189-201, 1997). As a result, the cauliflower mosaic virus 35S (CaMV 35S; Odell *et al.*, Nature 313 653-660, 1985) is currently the most widely used constitutive promoter in plant science. Its regulation is well known and it has been shown to work in many monocotyledonous and dicotyledonous plant species, and even in some fungal species. The main drawback of such a promoter however is its viral origin. Plants transformed with the CaMV35S promoter frequently show several undesirable characteristics, such as efficient silencing of the transgenes or increased recombination, leading to instability over time and over subsequent generations (Kohli *et al.*, Plant J. 17 591-601, 1999; Al-Kaff *et al.*, Nature Biotechnology 18 995-999, 2000). In addition, viral elements are often not desirable in transgenic products, both for regulatory reasons and for reasons of general public acceptance.

Rice *GOS2* cDNA was isolated and characterised in a screening program for genes that were expressed in many different rice tissues (de Pater *et al.*, Plant J. 2, 837-844, 1992). *GOS2* is a single copy gene that encodes an isoform of the translation initiation factor eIF1. Expression analysis revealed high mRNA levels in green shoots, etiolated shoots and in roots of seedlings. A similar expression pattern was detected in mature plants, though the expression was a bit lower in roots. High expression levels were also detected in plants or cell suspension cultures of another rice variety. An isolated promoter fragment was able to drive *GUS* expression in rice seedlings (leaf, root tissues and coleoptile), and in rice cell suspension cultures. The promoter was also active in maize, barley and ryegrass (de Pater *et al.*, Plant J. 2, 837-844, 1992; Hensgens *et al.*, Plant Mol. Biol. 22, 1101-1127, 1993). It has been shown to be active in other monocotyledonous plants, with expression levels similar to that of CaMV

35S. Barbour et al. (WO0020571) suggest, but did not show, the use of a GOS2 promoter from maize in dicotyledonous plants. There is however little homology between the maize GOS2 promoter and the rice GOS2 promoter that is the subject of the present invention.

5 It has now been found that a GOS2 regulatory sequence from rice is active in non-monocotyledonous plants, and has a similar expression pattern and strength to the CaMV 35S promoter. Consequently, the present invention concerns the use of a GOS2 regulatory sequence for driving constitutive expression of nucleic acids in non-monocotyledonous plants. The present invention therefore provides a good alternative for the CaMV 35S promoter. The
10 regulatory sequence of the present invention, when linked to a useful gene and transferred into a non-monocotyledonous plant or plant cell, enables expression of that useful gene. In addition, since the regulatory sequence of the present invention enables constitutive expression, it is possible to use it as a regulatory sequence to drive expression of the transferred useful gene in all tissues at any developmental stage. Moreover, as this regulatory
15 sequence originates from a monocotyledonous plant, it is less likely to be prone to silencing than dicotyledonous regulatory sequence or the CaMV35S promoter, when used to express a transferred useful gene in dicotyledonous plants, such as *Arabidopsis thaliana*.

Those skilled in the art will be aware that the invention described herein is subject to variations
20 and modifications other than those specifically described. It is to be understood that the invention described herein includes all such variations and modifications. The invention also includes all such steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any or more of these steps or features.

25 Throughout this specification, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not to the exclusion of any other integer or step or group of integers or steps.

30 As used herein, the term "derived from" shall be taken to indicate that a particular integer or group of integers has originated from the species specified, but has not necessarily been obtained directly from the specified source. This integer may be substantially modified from its native form in composition and/or environment through deliberate human manipulation.

35 In accordance with the present invention, there is provided the use of an isolated regulatory nucleic acid sequence comprising a regulatory sequence as represented by SEQ ID NO 1 or

by a functional fragment or functional variant thereof, for driving expression of an associated nucleic acid sequence in a non-monocotyledonous plant or plant cell.

5 The term "regulatory sequence" as used herein refer to DNA sequences which are necessary to effect expression of sequences to which they are ligated. The regulatory sequences may differ depending upon the intended host organism and upon the nature of the sequence to be expressed. The term "regulatory sequence" is intended to include, as a minimum, all components necessary for expression and optionally additional advantageous components. According to a preferred feature, the isolated regulatory nucleic acid sequence is a promoter
10 sequence. As used herein, a 'promoter' means a region of DNA upstream from the transcription start and which is involved in binding RNA polymerase and other proteins to start transcription. Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences derived from a classical eukaryotic genomic gene, including the TATA box which is required for accurate transcription initiation, with or
15 without a CCAAT box sequence and additional regulatory elements (i.e. upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner. Consequently, a repressible promoter's rate of transcription decreases in response to a repressing agent. An inducible promoter's rate of transcription increases in response to an inducing agent. A
20 constitutive promoter's rate of transcription is not specifically regulated, though it can vary under the influence of general metabolic conditions. The term "promoter" also includes the transcriptional regulatory sequences of a classical prokaryotic gene, in which case it may include a -35 box sequence and/or a -10 box transcriptional regulatory sequences. The term "promoter" is also used to describe a synthetic or fusion molecule, or derivative which confers,
25 activates or enhances expression of a nucleic acid molecule in a cell, tissue or organ.

The present invention thus also encompasses the use of fragments or variants of SEQ ID NO 1, which may be joined by any means known in the art, e.g., by ligation *in vitro*, to create hybrid regulatory sequences comprising mixtures of parts of regulatory elements from different
30 sources, either natural or synthetic. Therefore, the invention also provides a hybrid regulatory nucleic acid sequence comprising the regulatory sequence represented by SEQ ID NO 1 or a fragment or variant thereof, functionally combined with one or more other regulatory sequences or fragments thereof.

35 Regulatory sequences or promoters may contain additional copies of one or more specific regulatory elements, to further enhance expression and/or to alter the spatial expression and/or temporal expression of a nucleic acid molecule to which it is operably connected. Such

regulatory elements may be placed adjacent to a heterologous promoter sequence to drive expression of a nucleic acid molecule in response to e.g. copper, glucocorticoids, dexamethasone, tetracycline, gibberellin, cAMP, abscisic acid, auxin, wounding, ethylene, jasmonate or salicylic acid, or to confer expression of a nucleic acid molecule to specific cells, tissues or organs such as meristems, leaves, roots, embryo, flowers, seeds or fruits. In the context of the present invention, the regulatory sequence is active in non-monocotyledonous plants, and is represented by SEQ ID NO1, a functional fragment or a functional variant thereof. The term "functional", when used in respect of a nucleic acid or part thereof, is to be understood as having an essentially similar biological activity compared to the naturally occurring nucleic acid and being capable of driving expression in non-monocotyledonous plants.

"Expression" means the production of a protein or nucleotide sequence in a cell itself or in a cell-free system. It includes transcription into an RNA product, post-transcriptional modification and/or translation to a protein product or polypeptide from a DNA encoding that product, as well as possible post-translational modifications. A coding sequence can be transcribed in sense or antisense direction.

The term "nucleic acid sequence(s)", "nucleic acid molecule(s)", "polynucleotide(s)", "nucleotide sequence(s)", "DNA sequence(s)" or "gene(s)", when used herein refers to nucleotides, either ribonucleotides (RNA) or deoxyribonucleotides (DNA) or a combination of both, in a polymeric form of any length. These terms furthermore include double-stranded and single-stranded DNA and RNA. These terms also include nucleotide modifications known in the art, such as methylation, cyclisation, 'caps', substitution of one or more of the naturally occurring nucleotides with an analogue such as inosine, and polynucleotide backbone modifications. The term "isolated" in the present application means removed from its original environment. For example, a nucleic acid present in the natural state in an organism is not isolated, whereas the same nucleic acid separated from the adjacent nucleic acids in which it is naturally present, is regarded as being "isolated".

An associated nucleic acid sequence may comprise coding sequences or it may contain non-coding sequences. A "coding sequence" or "open reading frame" or "ORF" is defined as a nucleotide sequence that can be transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences, i.e. when the coding sequence or ORF is present in an expressible form. The coding sequence or ORF is bound by a 5' translation start codon and a 3' translation stop codon. A coding sequence or ORF may include, but is not limited to RNA, mRNA, cDNA, recombinant nucleotide sequences,

synthetically manufactured nucleotide sequences or genomic DNA. The coding sequence or ORF can be interrupted by intervening nucleic acid sequences.

5 Coding sequences, or genes essentially encoding the same protein but isolated from different sources, can consist of substantially divergent nucleic acid sequences. Reciprocally, substantially divergent nucleic acid sequences can be designed to effect expression of essentially the same protein. These nucleic acid sequences are the result of e.g. the existence of different alleles of a given gene, or of the degeneracy of the genetic code or of differences in codon usage. Thus, amino acids such as methionine and tryptophan are encoded by a single
10 codon whereas other amino acids such as arginine, leucine and serine can each be translated from up to six different codons. Preferred codon usage of various organisms can be found in <http://www.kazusa.or.jp/codon>. Allelic variants are further defined as comprising single nucleotide polymorphisms (SNPs) as well as small insertion/deletion polymorphisms (INDELs; the size of INDELs is usually less than 100 bp). SNPs and INDELs form the largest set of
15 sequence variants in naturally occurring polymorphic strains of most organisms.

The term "associated" as used herein, or "operably linked" refers to a juxtaposition between a regulatory and a coding sequence, wherein the components so described are in a relationship permitting them to function in their intended manner. A regulatory sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is
20 achieved under conditions compatible with the control sequences. In case the regulatory sequence is a promoter, it would be known to a skilled person that a double-stranded nucleic acid is preferable. The associated nucleic acid encompasses heterologous nucleic acids. Heterologous nucleic acids refer to nucleic acids derived from a separate genetic source, for
25 example nucleic acids that originate from within the cell but that are not naturally located in the cell, or that are located in a different chromosomal site of the cell. Heterologous nucleic acids may also be derived from other species and may be introduced as a transgene, for example, by transformation. This transgene may be substantially modified from its native form in composition and/or genomic environment through deliberate human manipulation. Also
30 expression of the native nucleic acid sequences may be modified by introduction in the plant of regulatory sequences according to the invention in such a way that the regulatory sequences are operably linked to the native nucleic acid. Native nucleic acid refer to nucleic acid in their natural location in the genome of an organism.

5 The term "fragment of a sequence" or "part of a sequence" means a truncated sequence of the original sequence in question. The truncated nucleic acid sequence can vary widely in length; the minimum size being a sequence of sufficient size to provide a sequence with at least a

comparable function and/or activity of the original sequence in question, while the maximum size is not critical. In some applications, the maximum size usually is not substantially greater than that required to provide the desired activity and/or function(s) of the original sequence. These terms are not restrictive to the content of the DNA fragment or segment, which can be any DNA, with any functionality. For example the DNA fragment or segments may consist of a promoter but may also comprise many genes, with or without additional control elements, it may just contain spacer sequences. A functional fragment of a promoter may be constructed by deleting parts at the 5' and/or 3' end, and/or by deleting internal parts and testing the remaining fragment for its ability to drive expression of a reporter gene. The expression level and pattern of this reporter gene construct can then be compared to that of the same reporter gene under control of the original promoter from which the fragment was derived.

Furthermore, a variant of a DNA fragment encompasses a sequence showing homology to the sequence referred to and is capable of promoting expression of an associated DNA sequence when reintroduced into a plant and hybridises to the sequence referred to or to a portion thereof under stringent conditions. The person skilled in the art will appreciate that in this case, reactions under stringent conditions for hybridisation are typically carried out at a temperature of between 60°C and 65°C in 0.3 strength citrate buffer saline containing 0.1% SDS followed by rinsing at the same temperature with 0.3 strength citrate buffer saline containing 0.1% SDS. For the purposes of defining the level of stringency, reference can conveniently be made to Sambrook et al. (2001) *Molecular Cloning: a laboratory manual*, 3rd Edition Cold Spring Harbor Laboratory Press, CSH, New York or to *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989).

A "non-monocotyledonous plant" comprises all plant species which belong to the kingdom Plantae as defined by ITIS [the Integrated Taxonomic Information System (<http://www.itis.usda.gov/index.html>)], including the class of Magnoliopsida but without the class of the Liliopsida (Liliatae, Monocotyledonae). The classes Magnoliopsida and Liliopsida are part of the division Magnoliophyta, which in turn belongs to the subkingdom Tracheobionta. The isolated nucleic acid according to the present invention can be used to drive gene expression in any non-monocotyledonous plant, in particular it is applicable to a dicotyledonous plant including a fodder or forage legume, ornamental plant, food crop, tree, or shrub. Preferred plant species include cotton, potato, tomato, cabbage, sugar beet, soybean, bean, sunflower, peas.

Advantageously, the regulatory sequence represented by SEQ ID NO 1 or a functional fragment or functional variant thereof is capable of driving expression of an associated nucleic

acid sequence in a non-monocotyledonous plant. Such an associated nucleic acid may be any heterologous DNA sequence derived from any organism which sequence is to be expressed in a non-monocotyledonous plant. The heterologous DNA sequence may be obtained from any organism, including plants, which plant species may be the same plant species into which the sequence is to be introduced, or it may be different. Additionally or alternatively, the associated nucleic acid sequence may be an endogenous nucleic acid operably linked to the regulatory sequence according to the invention.

Therefore, the invention also provides use of an isolated regulatory nucleic acid as defined above for driving expression of an associated nucleic acid in a non-monocotyledonous plant or plant cell, wherein said associated nucleic acid is an isolated nucleic acid or a nucleic acid endogenous to the host cell in which said isolated regulatory nucleic acid sequence is introduced.

The recombinant nucleic acid encompasses any means for the cloning of and/or transfer of a nucleic acid into a host cell, including "vectors" or "vector sequences", which may be introduced in an organism by transformation or transfection, and which may either be integrated into the genome of the host cell or maintained in some form extrachromosomally. Vector sequences generally comprise a set of unique sites recognised by restriction enzymes, the multiple cloning site (MCS), wherein one or more non-vector sequence(s) can be inserted.

With "non-vector sequence" or "insert" is accordingly meant the desired DNA segment which one wishes to clone in one or more of the sites of the MCS, comprised within a vector. The insert can have one or more genes.

"Expression vectors" form a subset of vectors, which, by virtue of comprising the appropriate regulatory sequences enabling the creation of an expressible format for the inserted non-vector sequence(s), thus allowing transcription and/or translation of a nucleic acid inserted therein. Expression vectors are known in the art and enable protein expression in organisms including bacteria (e.g. *Escherichia coli*), fungi (e.g. *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pichia pastoris*), insect cells (e.g. baculoviral expression vectors), animal cells (e.g. COS or CHO cells) and plant cells (e.g. potato virus X-based expression vectors, see e.g. Vance *et al.* 1998 - WO9844097). Expression vectors can for instance be cloning vectors, binary vectors or integrating vectors.

Regulatory sequences ensuring expression in prokaryotic and/or eukaryotic cells, particularly in non-monocotyledonous plants, are well known to those skilled in the art. A preferred

regulatory sequence is the regulatory sequence listed in SEQ ID NO 1, a functional fragment or a functional variant thereof. When used in eukaryotic cells vectors may also comprise terminators which contain polyadenylation signals ensuring 3' processing and polyadenylation of a primary transcript and termination of transcription. In plant cells the termination signals usually employed are from the *nopaline synthase* gene or the CaMV 35S terminator. Additional regulatory elements may include transcriptional as well as translational enhancers. A plant translational enhancer often used is the CaMV omega sequence. The inclusion of an intron has been shown to increase expression levels by up to 100-fold in certain plants (Maiti *et al.*, Transgenic Research 6 143-156, 1997; Ni, Plant Journal 7 (1995), 661-676).

Advantageously, vectors used in the invention comprise a selectable and/or scorable marker. Selectable marker genes useful for the selection of transformed plant cells, callus, plant tissue and plants are well known to those skilled in the art. Antimetabolite resistance is very useful as a basis for selection: for example, the *dhfr* gene confers resistance to methotrexate (Reiss *et al.*, Plant Physiol. (Life Sci. Adv.) 13 143-149, 1994); the *npt* gene confers resistance to the aminoglycosides neomycin, kanamycin and paromomycin (Herrera-Estrella *et al.*, EMBO J. 2 987-995, 1983); or the *hpt* gene, which confers resistance to hygromycin (Marsh, Gene 32 481-485, 1984). Additional selectable markers genes have been described, such as *trpB*, which allows cells to utilize indole in place of tryptophan; *hisD*, allowing cells to utilize histinol in place of histidine (Hartman, Proc. Natl. Acad. Sci. USA 85 8047, 1988); *mannose-6-phosphate isomerase* enabling cells to utilize mannose (WO 94/20627) and *ornithine decarboxylase* which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine or DFMO (McConlogue, 1987, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed.) or the *deaminase* gene from *Aspergillus terreus* which confers resistance to Blasticidin S (Tamura, Biosci. Biotechnol. Biochem. 59 2336-2338, 1995). Useful scorable markers are also known to those skilled in the art and are commercially available. Advantageously, the marker is a gene encoding luciferase (Giacomin, Pl. Sci. 116 59-72, 1996; Srikantha, J. Bact. 178 121, 1996), green fluorescent protein (Gerdes, FEBS Lett. 389 44-47, 1996; Haseloff *et al.*, Proc. Natl. Acad. Sci U.S.A. 94 2122-2127, 1997) or β -glucuronidase (Jefferson, EMBO J. 6 3901-3907, 1987). Vectors may further provide primer sites, e.g., for PCR, transcriptional and/or translational initiation and/or regulation sites, recombinational sites, replicons, etc.

The current invention clearly includes any recombinant nucleic acid, vector or expression vector comprising the regulatory sequence according to the present invention and/or a non-vector sequence as defined supra.

According to another embodiment, the invention relates to a non-monocotyledonous plant cell comprising a recombinant nucleic acid as described above. The recombinant nucleic acid may be stably integrated into the genome of the non-monocotyledonous plant cell. The invention also provides a plant cell culture, callus, a plant or a plant part derived from these plant cells.

5 Also within the scope of the present invention is a harvestable part, organ, tissue or propagation material of a plant according to the invention, comprising a nucleic acid sequence comprising a regulatory sequence represented by SEQ ID NO: 1 or a fragment or variant thereof.

10 Furthermore, there is provided a method for expression of a nucleic acid sequence in a non-monocotyledonous plant cell, which method comprises introducing into this plant cell a regulatory sequence represented by SEQ ID NO 1 or a functional fragment or functional variant thereof, wherein the regulatory sequence is capable of driving expression of the nucleic acid sequence which is either an isolated or endogenous nucleic acid sequence.

15 Means for introducing recombinant DNA into plant tissue or cells include, but are not limited to, transformation using CaCl_2 and variations thereof (Hanahan, J. Mol. Biol. 166 (4) 557-580, 1983)), direct DNA uptake into protoplasts (Krens *et al.*, Nature 296 72-74, 1982; Paszkowski *et al.*, EMBO J. 3 2717-2722, 1984), PEG-mediated uptake to protoplasts (Armstrong *et al.*,
20 Plant Cell Reports 9 335-339, 1990) microparticle bombardment, electroporation (Fromm *et al.*, Proc. Natl. Acad. Sci. U.S.A. 82 5824-5826, 1985) microinjection of DNA (Crossway *et al.*, Mol. Gen. Gen. 202 179-185, 1986; Fromm *et al.* Proc. Natl. Acad. Sci. U.S.A. 82 5824-5826, 1985), microparticle bombardment of tissue explants or cells (Christou *et al.*, Plant Physiol. 87 671-674, 1988), vacuum-infiltration of tissue with nucleic acid, or in the case of plants, T-DNA-
25 mediated transfer from *Agrobacterium* to the plant tissue as described essentially (An *et al.*, EMBO J. 4 277-284, 1985; Dodds, Plant genetic engineering, 1985; Herrera-Estrella *et al.*, EMBO J. 2 987-995, 1983; Herrera-Estrella *et al.*, Nature 303 209-213, 1983).

30 For microparticle bombardment of cells, a microparticle is propelled into a cell to produce a transformed cell. Any suitable ballistic cell transformation methodology and apparatus can be used in performing the present invention. Exemplary apparatus and procedures are disclosed by Stomp *et al.* (U.S. Patent No. 5122466) and Sanford and Wolf (U.S. Patent No. 4945050). When using ballistic transformation procedures, the gene construct may incorporate a plasmid capable of replicating in the cell to be transformed. Examples of microparticles suitable for use
35 in such systems include 1 to 5 μm gold spheres. The DNA construct may be deposited on the microparticle by any suitable technique, such as by precipitation.

The nucleic acid as defined above in the present invention, or a genetic construct comprising it, may be introduced into a cell using any known method for the transfection or transformation.

5 The invention further relates to a method for the production of non-monocotyledonous transgenic plants, plant cells or plant tissues comprising the introduction in a non-monocotyledonous plant, plant cell or plant tissue of a first nucleic acid molecule according to the invention operably linked to an associated nucleic acid sequence, wherein the associated nucleic acid sequence is heterologous to the first nucleic acid. According to another preferred embodiment, the invention relates to a non-monocotyledonous transgenic plant cell or plant as
10 described here above wherein the nucleic acid of the invention is stably integrated into the genome of the non-monocotyledonous plant cell or plant.

Upon transformation of a cell with the genetic construct of the invention, a whole plant may be regenerated therefrom. Plant tissue capable of subsequent clonal propagation, whether by
15 organogenesis or embryogenesis, may be transformed with a gene construct of the present invention and a whole plant regenerated therefrom. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g.,
20 apical meristem, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem). The term "organogenesis", as used herein, means a process by which shoots and roots are developed sequentially from meristematic centres. The term "embryogenesis", as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or
25 gametes.

The generated transformed non-monocotyledonous plants may be propagated by a variety of means, such as by clonal propagation or classical breeding techniques. For example, a first generation (or T1) transformed plant may be selfed to give homozygous second generation (or
30 T2) transformants, and the T2 plants further propagated through classical breeding techniques. The generated transformed plants contemplated herein may take a variety of forms. For example, they may be chimeras of transformed cells and non-transformed cells; clonal transformants (e.g., all cells transformed to contain the expression cassette); grafts of transformed and untransformed tissues (e.g., in plants, a transformed rootstock grafted to an
35 untransformed scion).

The invention not only relates to a transgenic non-monocotyledonous plant or plant tissue comprising plant cells as herein described but extends also to the harvestable parts of the transgenic plant, preferably selected from the group consisting of seeds, leaves, flowers, fruits, stem cultures, roots, tubers, rhizomes and bulbs. The invention also relates to the progeny
5 derived from any of these transgenic plants or plant parts.

The present invention also encompasses use of an isolated nucleic acid sequence comprising a regulatory sequence as represented by SEQ ID NO 1 or a functional fragment or functional variant thereof, characterised as above for driving constitutive expression of an associated
10 sequence in a non-monocotyledonous plant cell or plant. Also included is the use of a hybrid regulatory nucleic acid as defined above for driving expression of an associated nucleic acid sequence in a non-monocotyledonous plant cell or plant.

Description of figures

- 15 **Fig 1:** Typical GUS expression pattern in an *Arabidopsis thaliana* plant transformed with a p2203 construct and a control plant. The GOS2 regulatory sequence shows a strong constitutive expression in shoot tissues, including the petioles, the vascular system, trichomes and in root tips. Expression in other root tissues is lower, but still easily noticeable.
- 20 **Fig 2:** GUS expression under control of the GOS2 regulatory sequence (lines AT0476-09 and AT0476-11), compared to GUS expression under control of the 35S promoter (line WS35S:GUS). For each line, 5 replicas were tested. The plants shown have 2 cotyledons and 2 true leaves and were stained for 1 hour (panel a) or were stained overnight (panel b).
- 25 **Fig 3:** GUS expression under control of the GOS2 regulatory sequence (lines AT0476-09 and AT0476-11), compared to GUS expression under control of the 35S promoter (line WS35S:GUS). For each line, 5 replicas were tested. The plants shown are in the 10 leaf stage and were stained for 1 hour (panel a) or were stained overnight (panel b).
- 30 **Fig 4:** GUS expression under control of the GOS2 regulatory sequence (lines AT0476-09 and AT0476-11), compared to GUS expression under control of the 35S promoter in line WS35S:GUS. For each line, 5 replicas were tested. The plants shown are in the adult stage with an inflorescence and were stained for 1 hour (panel a) or were stained overnight (panel b).

35

Examples

The present invention will now be illustrated with the following examples.

DNA manipulation

- 5 All DNA procedures were performed according to standard protocols (Maniatis T *et al.* (2001). Molecular Cloning: a laboratory manual, Cold Spring Harbor Laboratory Press, CSH, New York).

Example 1: Cloning of GUS with intron coding sequence

- 10 The *E. coli* β -glucuronidase gene, interrupted by the second intron of the Potato light-inducible tissue-specific ST-LS1 gene, was amplified from vector pTHW136 (with a cassette identical to the one of pMOG553, MOGEN international n.v, LEIDEN, NETHERLANDS). The cDNA was amplified by PCR with Platinum Pfx DNA polymerase (Invitrogen) and using a sense primer including attB1 (SEQ ID NO2) and an antisense primer including attB2 (SEQ ID NO3).

- 15 Conditions for PCR were: 2 minutes of denaturation at 94°C (1 cycle); 35 cycles of 1 minute denaturation at 94°C, 1 minute annealing at 58°C and 2 minutes amplification at 68°C, and finally 1 cycle of 5 minutes elongation at 68°C. A prominent fragment with the expected size of 2 kb was isolated from gel and purified using the Zymoclean Gel DNA Recovery Kit (Zymo
20 Research, Orange, California).

- The purified PCR fragment was used in a standard Gateway™ BP reaction (Invitrogen) with pDONR201 as a recipient vector. The identity and base pair composition of the insert was confirmed by sequencing. The integrity of the resulting plasmid was verified using restriction
25 digests and was given the designation p0604 (entry clone). pDONR201 was obtained from Invitrogen.

- p0604 is a Gateway™ "entry clone", and was used as such in a standard Gateway™ LR reaction, with p0640 as "destination vector". The resulting vector was p02203, its integrity was
30 also checked by restriction digest analysis. The vector, containing as functional elements within the T-DNA region a selectable marker gene and a "Gateway cassette" intended for LR cloning of sequences of interest, was used for the transformation of *Arabidopsis thaliana*. Expression of these sequences of interest, upon being recombined into p0640, was driven by the regulatory sequence as listed in SEQ ID NO1.

35

Example 2: Transformation of plants

Cultivation of the parental plants

For the parental plants, approximately 12mg of wild-type *Arabidopsis thaliana* (ecotype Columbia) seeds were suspended in 27.5 ml of 0.2 % agar solution. The seeds were
5 incubated for 2 to 3 days at a temperature of 4°C and then sown. The plants were germinated under standardised conditions: 22°C during the day, 18°C at night, with a relative humidity of 65-70%, 12 hours of photoperiod and sub-irrigation with water for 15 min every 2 or 3 days. The seedlings were then transplanted into pots with a diameter of 5.5cm, containing a mixture of sand and peat in a ratio of 1 to 3. The plants were then further grown under the same
10 standard conditions as mentioned above.

Agrobacterium growth conditions and preparation

Agrobacterium strain C58C1RIF with helper plasmid pMP90 and vector p2203 was inoculated in a 50 ml plastic tube containing 1 ml LB (Luria-Bertani Broth) without antibiotic. The culture
15 was shaken for 8-9h at 28°C. Next, 10 ml of LB without antibiotic was added to the plastic tube and shaken overnight at 28°C. Following this, the OD at 600 nm was measured. At an optical density of approximately 2.0, 40 ml of 10% sucrose and 0.05% Silwet L-77 (a mixture of 84 % polyalkyleneoxide modified heptamethyltrisiloxane and 16% allyloxypolyethyleneglycol methyl ether, OSI Specialties Inc) was added to the culture. The *Agrobacterium* culture thus
20 obtained was used to transform the grown plants.

Flower dip

When each parental flower had one inflorescence of 7-10 cm in height, the inflorescences were submerged in the *Agrobacterium* culture and swirled gently for 2-3 seconds. For each
25 transformation, 2 plants were used. Hereafter, the plants were returned to the growing conditions as described above.

Seed collection

5 weeks after the flowers were dipped in the *Agrobacterium* culture, watering of the plants was
30 stopped. The plants were further incubated at 25°C with a photoperiod of 20 hours. One week later, the seeds were harvested and placed in a seed drier during one week. The seeds were then cleaned, collected in 15 ml plastic tubes and stored at 4°C until further processing.

T1 seeds generation

35 Transgenic T0 seeds were selected for their marker expression, and germinated. Plants were grown as described above, until T1 seeds were harvested. The different lines were given the name AT0476-xx, where xx is a number.

Example 3: Comparison of the GOS2 regulatory sequence and the 35S promoter in Arabidopsis:

The expression pattern of the β -glucuronidase gene and strength of the GOS2 regulatory sequence were compared with those of the 35S promoter in *Arabidopsis*.

Plant material used:

- AT0476-11: cassette GOS2-GUS and screenable marker cassette (vector p2203) (T1 seeds)
- AT0476-09: cassette GOS2-GUS and screenable marker cassette (vector p2203) (T1 seeds)
- WS35S:GUS: cassette 35S-GUS, no marker selection, homozygous line from Versailles (T2 homozygous seeds), Wassilewskija (WS) background.

Procedure:

Around 100 seeds, after being kept at a temperature of 4°C for at least 2 days, were sown in soil and the plants were cultivated under standard conditions: 22°C during the day, 18°C at night, 65-70% relative humidity, 12 hours of photoperiod, sub-irrigation with water for 15 min every 2 or 3 days.

When the plants had 2 cotyledons and 2 true leaves, marker expression in AT0476 lines was monitored and plants with a uniform marker expression pattern were retained.

Next, 10 plants of each GOS2 or 35S lines were carefully isolated from the soil and labelled for identification. 5 of the plants were stained for GUS during 1 hour, the other 5 plants were stained overnight. The short staining time was needed to assess qualitatively the relative strength of the promoters. If the staining reaction was saturated in less than 1 hour, the incubation time is to be reduced. In case the staining was very faint blue after one hour, the incubation time may be increased to 2-3 hours. Finally, the stained plants were photographed. For each line, 20 other plants were carefully isolated from the soil and transferred in individual small pots for further cultivation. When the plants had reached the 10-leaf stage, again 10 plants of each line were isolated, labelled and stained as described above. The stained plants were photographed. At flowering time, 10 plants per line with siliques in various maturation stages were isolated and processed as above.

Plant GUS staining procedure

The material was covered with 90 % ice-cold acetone and incubated for 30 min at 4°C. After 3 washes of 5 minutes with Tris buffer [15,76 g Trizma HCl (Sigma T3253) + 2,922 g NaCl in 1 l double distilled water, adjusted pH to 7,0 with NaOH], the material was immersed in a Tris/ferricyanate/X-Gluc solution [9,8 ml Tris buffer + 0,2 ml ferricyanate stock (0,33 g

Potassium ferricyanate (Sigma P3667) in 10 ml Tris buffer)+ 0,2 ml X-Gluc stock (26,1 mg X-Gluc (Europa Bioproducts ML 113A) in 500 μ l DMSO)]. Vacuum infiltration was applied for 15 to 30 minutes. Next, the samples were incubated for an appropriate time (up to 16 hours) at 37 °C for development of the blue colour. The samples were washed 3 times during 5 minutes with Tris buffer. Chlorophyll was extracted in a series of 50 %, 70 % and 90 % ethanol (each for 30 minutes) with refreshments of the 90 % solution if necessary.

Results

The GOS2 regulatory sequence shows a strong constitutive expression in shoot tissues, including the petioles, the vascular system and trichomes. Except for the root tips, the expression in root tissues is lower, but is still easily detectable (Fig 1).


For all the plants with the same incubation time during staining (1hr or overnight), the pattern of GUS expression was similar. The staining of GOS2 plants was as strong as for 35S plants, at all stages that were sampled (2 leaves+ 2 cotyledons, 10 leaf stage and flowering stage). GOS2 expression appears equally strong and equally constitutive as 35S expression in *Arabidopsis* (Figs 2, 3 and 4).

Claims

- 1) Use of an isolated regulatory nucleic acid sequence comprising a regulatory sequence as represented in SEQ ID NO 1 or a functional fragment or a functional variant thereof, for driving expression of an associated nucleic acid sequence in a non-monocotyledonous plant or plant cell.
- 2) Use of an isolated regulatory nucleic acid sequence according to claim 1, wherein said associated nucleic acid sequence is an isolated nucleic acid sequence or a nucleic acid sequence endogenous to the host cell in which said isolated regulatory nucleic acid sequence is introduced.
- 3) A non-monocotyledonous plant cell comprising or having stably integrated into its genome a recombinant nucleic acid as represented in SEQ ID NO 1 or a functional fragment or a functional variant thereof.
- 4) A non-monocotyledonous plant cell according to claim 3, wherein said non-monocotyledonous plant cell is derived from a fodder or forage legume, ornamental plant, food crop, tree or shrub, preferably from cotton, potato, tomato, cabbage, sugar beet, soybean, bean, sunflower or peas.
- 5) A plant cell culture, callus or a plant consisting essentially or in part of plant cells according to claim 3 or 4.
- 6) A harvestable part, organ, tissue or propagation material of a plant according to claim 5.
- 7) Method for expression of a nucleic acid sequence in a non-monocotyledonous plant or plant cell, said method comprising introducing into said plant or plant cell a regulatory sequence represented by SEQ ID NO 1 or a functional fragment or functional variant thereof, wherein said regulatory sequence is capable of driving expression of said nucleic acid sequence which is either an isolated or an endogenous nucleic acid sequence.

1/5

A



B




FIGURE 1

2/5



A 1h staining**B** ON staining

FIGURE 2

3/5




FIGURE 3

4/5




FIGURE 4

5/5



FIGURE 4 (continued)

SEQUENCE LISTING

<110> CropDesign N.V.

<120> Regulatory sequence

<130> CD-072-PCT

<150> EP 03075207.5

<151> 2003-01-21

<160> 3

<170> PatentIn version 3.1

<210> 1

<211> 2195

<212> DNA

<213> Oryza sativa

<400> 1

aatccgaaaa	gtttctgcac	cgttttcacg	tcctaactaa	caatataggg	aacgtgtgct	60
aaatataaaa	tgagacctta	tatatgtagc	gctgataact	agaactatgt	aagaaaaact	120
catccaccta	cttttagtggc	aatcgggcta	aataaaaaag	agtcgctaca	ctagtttcgt	180
tttccttagt	aattaagtgg	gaaaatgaaa	tcattattgc	ttagaatata	cgttcacatc	240
tctgtcatga	agttaaatta	ttcgaggtag	ccataattgt	catcaaactc	ttcttgaata	300
aaaaaatctt	tctagctgaa	ctcaatgggt	aaagagagat	atataattat	ataattttat	360
agaatgaaga	tattctgaac	gtatcggcaa	agattttaa	atataattat	ataattttat	420
agtttgtgca	ttcgttatat	cgcacgtcat	taaggacatg	tcttactcca	tctcaatttt	480
tatttagtaa	ttaaagacaa	ttgacttatt	tttattattt	atcttttttc	gattagatgc	540
aaggtagtta	cgcacacact	ttgtgctcat	gtgcatgtgt	gagtgacact	cctcatcac	600
gttcaactag	cgacacatct	ccaatatcac	tcgcctat	aatacattta	ggtagcaata	660
tctgaattca	agcacttcac	catcaccaga	ccacttttaa	taatatctaa	aatacaaaaa	720
ataattttac	agaatagcat	gaaaagtatg	aaacgaacta	tttaggtttt	tcacatacaa	780
aaaaaaaaag	aattttgtct	gtgcgcgagc	gccaatctcc	catattgggc	acacaggcaa	840
caacagagtg	gctgcccaca	gaacaaccca	caaaaaacga	tgatctaacg	gaggacagca	900
agtcgcgaac	aaccttttaa	cagcaggctt	tgccggccagg	agagaggagg	agaggcaaag	960
aaaaccaagc	atcctcctcc	tcccatctat	aaattcctcc	ccccttttcc	cctctctata	1020
taggaggcat	ccaagccaag	aagagggaga	gcaccaagga	cacgcgacta	gcagaagccg	1080
agcgaccgcc	ttcttcgatc	catatcttcc	ggctcgagttc	ttggctgatc	tcttccctcc	1140
tccacctcct	cctcacaggg	tatgtgccct	tcgggtgttc	ttggatttat	tggtctaggt	1200
tgtgtagtac	ggcgttgat	gttaggaaag	gggatctgta	tctgtgatga	ttcctgttct	1260
tggatttggg	atagagggt	tcttgatgtt	gcattgttatc	ggttcgggtt	gattagtagt	1320
atggttttca	atcgtctgga	gagctctatg	gaaatgaaat	ggtttagggg	acggaatctt	1380
gcgattttgt	gagtagcttt	tggttgaggt	aaaatcagag	caccgggtgat	tttgcttggt	1440
gtaataaaaag	tacatttggt	tggtcctcga	ttctggtagt	gatgcttctc	gatttgacga	1500
agctatcctt	tgtttattcc	ctattgaaca	aaaataatcc	aaotttgaag	acggtcccgt	1560
tgatgagatt	gaatgattga	ttcttaagcc	tgtccaaaat	ttcgcagctg	gcttggttag	1620
atacagtagt	ccccatcacg	aaattcatga	aaacagttat	aatcctcagg	aacaggggat	1680
tccctgttct	tcgatttgc	tttagtccca	gaattttttt	tcccaaatat	cttaaaaagt	1740
cactttctgg	ttcagttcaa	tgaattgatt	gctacaaata	atgcttttat	agcgttatcc	1800
tagctgtagt	tcagtttata	gtaataatcc	ctatagttta	gtcaggagaa	gaacttatcc	1860
gatttctgat	ctccattttt	aatttatatga	aatgaactgt	agcataagca	gtattcat	1920
ggattatttt	ttttattagc	tttcacccct	tcattattct	gagctgaaag	tctggcatga	1980
actgtcctca	atcttgtttt	caaattcaca	tcgattatct	atcgattatc	ctcttgatc	2040
tacctgtaga	agtttctttt	tgggttattcc	ttgactgctt	gattacagaa	agaaatttat	2100
gaagctgtaa	tcgggatagt	tatactgctt	gttcttatga	ttcatttcc	ttgtgcagtt	2160
cttggtgtag	cttgccactt	tcaccagcaa	agttc			2195

CD0723PCT.ST25.txt

<210> 2
<211> 61
<212> DNA
<213> Artificial sequence

<220>
<223> sense primer attB1

<400> 2
ggggacaagt ttgtacaaaa aagcagggtt cacaatgtta cgtcctgtag aaacccaac 60
c 61

<210> 3
<211> 52
<212> DNA
<213> Artificial sequence

<220>
<223> antisense primer attB2

<400> 3
ggggaccact ttgtacaaga aagctgggtt tggtgattca ttgtttgcct cc 52

(12) INTERNATIONAL PUBLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
5 August 2004 (05.08.2004)

PCT

(10) International Publication Number
WO 2004/065596 A3

(51) International Patent Classification: C12N 15/82

(21) International Application Number:
PCT/EP2004/000645

(22) International Filing Date: 21 January 2004 (21.01.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03075207.5 21 January 2003 (21.01.2003) EP(71) Applicant (for all designated States except US):
CROPDESIGN N.V. [BE/BE]; Technologiepark 3,
Zwijnaarde 9052 (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HATZFELD, Yves
[FR/FR]; 186 rue des Postes, Lille 59000 (FR). INZE, Dirk
[BE/BE]; Driesstraat 18, Moorsel- Aalst 9310 (BE).(74) Common Representative: CROPDESIGN N.V.; Tech-
nologiepark 3, Zwijnaarde 9052 (BE).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
30 September 2004For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.(54) Title: USE OF THE REGULATORY SEQUENCE OF THE RICE GOS2 GENE FOR THE GENE EXPRESSION IN DI-
COTYLEDONOUS PLANTS OR PLANT CELLS(57) Abstract: The present invention relates to the field of plant molecular biology. In particular, it describes the use of a regu-
latory nucleic acid sequence of the rice gene GOS2 for the regulation of gene expression in plant cells derived from plants other
than monocotyledonous plants. The use of the regulatory sequence of the present invention results in constitutive expression with
expression levels similar to that of CaMV 35S. The present invention also relates to vectors and host cells comprising these nucleic
acid sequences. The invention further relates to transgenic cells and plants comprising these sequences and to methods for obtaining
such cells and plants.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/000645

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/82

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, BIOSIS, EMBASE, MEDLINE, PAJ, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATER DE B S ET AL: "THE PROMOTER OF THE RICE GENE GOS2 IS ACTIVE IN VARIOUS DIFFERENT MONOCOT TISSUES AND BINDS RICE NUCLEAR FACTOR ASF-1" PLANT JOURNAL, BLACKWELL SCIENTIFIC PUBLICATIONS, OXFORD, GB, vol. 2, no. 6, 1992, pages 837-844, XP000907326 ISSN: 0960-7412 cited in the application the whole document	1-7
A	WO 92/20807 A (SALK INST FOR BIOLOGICAL STUDI) 26 November 1992 (1992-11-26) the whole document ----- -/--	1-7

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

29 July 2004

Date of mailing of the international search report

19/08/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Grötzinger, T

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/000645

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>OUELLET F VAZQUEZ-TELLO A SARHAN F: "The wheat wcs120 promoter is cold-inducible in both monocotyledonous and dicotyledonous species"</p> <p>FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 423, no. 3, 27 February 1998 (1998-02-27), pages 324-328, XP004261912 ISSN: 0014-5793 the whole document</p>	1-7
A	<p>LUAN S ET AL: "A RICE CAB GENE PROMOTER CONTAINS SEPARATE CIS-ACTING ELEMENTS THAT REGULATE EXPRESSION IN DICOT AND MONOCOT PLANTS"</p> <p>PLANT CELL, vol. 4, no. 8, 1992, pages 971-981, XP002283900 ISSN: 1040-4651 the whole document</p>	1-7
A	<p>HAUPTMANN R M ET AL: "PROMOTER STRENGTH COMPARISONS OF MAIZE SHRUNKEN 1 AND ALCOHOL DEHYDROGENASE 1 AND 2 PROMOTERS IN MONOCOTYLEDONOUS AND DICOTYLEDONOUS SPECIES"</p> <p>PLANT PHYSIOLOGY (ROCKVILLE), vol. 88, no. 4, 1988, pages 1063-1066, XP001182033 ISSN: 0032-0889 the whole document</p>	1-7
A	<p>WO 01/27296 A (MATIN HELLER TERESA ; UNITED KINGDOM GOVERNMENT (CA); ALLARD SHARON) 19 April 2001 (2001-04-19) the whole document</p>	1-7
A	<p>DATABASE BIOSIS DENG, W.; SHAO, Q.; JIANG, X.: "Expression of a zein gene Z4 introduced into dicotyledonous Solanum nigrum" XP002283901 Database accession no. PREV198987070807 abstract</p>	1-7

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2004/000645

Box No. I **Nucleotide and/or amino acid sequence(s) (Continuation of Item 1.b of the first sheet)**

1. With regard to any nucleotide and/or amino acid sequence disclosed in the International application and necessary to the claimed invention, the International search was carried out on the basis of:
- a. type of material
- ☒ a sequence listing
- ☐ table(s) related to the sequence listing
- b. format of material
- ☒ in written format
- ☒ in computer readable form
- c. time of filing/furnishing
- ☒ contained in the international application as filed
- ☒ filed together with the international application in computer readable form
- ☐ furnished subsequently to this Authority for the purpose of search
2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/000645

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9220807	A	26-11-1992	US 5399680 A	21-03-1995
			CA 2108549 A1	23-11-1992
			EP 0585375 A1	09-03-1994
			JP 6510662 T	01-12-1994
			WO 9220807 A1	26-11-1992
			US 5695939 A	09-12-1997
WO 0127296	A	19-04-2001	US 6326528 B1	04-12-2001
			AU 7766000 A	23-04-2001
			WO 0127296 A1	19-04-2001
			CA 2387440 A1	19-04-2001